

JCS53 U.S. PTO
12/22/99

12-27-99

R

Attorney Docket No.: _____

Patent Application


JCS53 U.S. PTO
09/470603
12/22/99

CERTIFICATE OF EXPRESS MAILING

"Express Mail" mailing label number: EE683174198US
Date of Deposit: December 22, 1999

I hereby certify that this paper or fee is being deposited with the United State Postal Service "Express Mail Post Office to Addressee" service under 37 CFR §1.10 on the date indicated above and is addressed to: BOX APPLICATION, Assistant Commissioner for Patents, Washington, DC 20231

Peter J. Manso December 22, 1999
Name of person mailing paper of fee Date


Signature

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re patent application of: Dave Bova
For: **NICOTINIC ACID COMPOSITIONS FOR TREATING
HYPERLIPIDEMIA AND RELATED METHODS THEREFOR**
Examiner:
Art Unit:
Express Mail Label No.: EE683174198US

BOX APPLICATION
Assistant Commissioner
for Patents
Washington, D.C. 20231

Sir:

PATENT APPLICATION TRANSMITTAL LETTER

Transmitted herewith for filing as a continuation patent application under 35U.S.C. §111(a) and §120 and under 37 C.F.R. §1.53(b) and §1.78, please find the following:

- X Specification, claims and abstract of the above-referenced patent application (total of 32 pages). I hereby verify that the attached papers are a true copy of what is shown in my records to be the above-identified prior application, including the oath or declaration originally filed.

_____ sheet(s) of drawing(s) (____ formal/ ____ informal)

☒ Combined Declaration and Power of Attorney (previously filed)

☒ The prior grandparent and patent applications are assigned of record to KOS Pharmaceuticals, Inc.

☒ A verified statement that this filing is by a small entity has been filed in the grandparent and parent applications and such status is still proper and desired (37 C.F.A. §1.9 and 1.27).

☒ Preliminary Amendment is enclosed.

☒ Return Post Card is enclosed.

The filing fee has been calculated as shown below:

	NO FILED	NO. EXTRA	RATE	FEE
BASIC FEE				\$760
TOTAL CLAIMS	12-20	0	\$18	\$0
INDEPENDENT CLAIMS	1-3	0	\$78	\$0
MULTIPLE DEPENDENT CLAIM(S) PRESENTED			\$260	\$0
TOTAL FEES:				\$760
Deduct one-half of fee for Small Entity				-380
TOTAL AMOUNT DUE:				\$380

☒ Please charge my Deposit Account No. 50-0951/20720-95328 (KOS) in the amount of \$380. A copy of this request is submitted herewith.

_____ A check in the amount of \$_____ is attached. Please charge any deficiency or credit any overpayment to Deposit Account No. _____.

☒ The Commissioner is hereby authorized to charge payment of the following fees or other deficient fees owed during the pendency of this application or credit any overpayment to Deposit Account No. 50-0951.

- ☒ Any patent application processing fees under 37 CFR 1.17 and under 37 CFR 1.20(d).
- ☒ The issue fee set in §37 CFR 1.18 at or before mailing of the Notice of Allowance, pursuant to 37 CFR 1.311(b).
- ☒ Any filing fees under §37 CFR 1.16 including all extension of time fees that are due pursuant to 37 CFR 1.316(a) during the pendency of this application.
- ☒ With respect to the prior copending U.S. patent applications from which this application claims benefit under 35 U.S.C. §120, the inventor(s) in this application is (are) the same.
- ☒ The inventorship for all claims in this application are the same.
- ☒ The Power of Attorney in the prior parent application is to Peter J. Manso, Registration No. 32,264.
- ☒ Address all future communications to:

Peter J. Manso
Akerman Senterfitt & Eidson, P.A.
Las Olas Centre II, Suite 1600
350 East Las Olas Blvd.
Ft. Lauderdale, FL 33301
Telephone: 954-463-2700
Facsimile: 954-463-2224

- ☒ A notification of the filing of this continuation is being filed in the parent application from which this application claims priority under 35 U.S.C. §120.

Pursuant to the provisions of 37 C.F.R. §1.10 and 1.53, applicant requests that this application be assigned a serial number and a filing date of December 22, 1999, the date upon which this application for U.S. patent was mailed to the U.S. Patent and Trademark Office by Express Mail.


I hereby declare further that all statement made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statement and the like so made are

punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statement may jeopardize the validity of the application or any patent issuing thereon.

Date: December 22, 1999

Respectfully Submitted,

By and For
AKERMAN SENTERFITT & EIDSON, P.A.



Peter J. Manso
Registration No. 32264
Attorney or Agent of Record

Akerman, Senterfitt & Eidson, P.A.
Las Olas Centre II, Suite 1600
350 East Las Olas Boulevard
Fort Lauderdale, Florida 33301
Telephone: 954-463-2700
Facsimile: 954-463-2224

Attorney Docket No.: _____

Patent Application

CERTIFICATE OF EXPRESS MAILING

"Express Mail" mailing label number: EE683174198US
Date of Deposit: December 22, 1999

I hereby certify that this paper or fee is being deposited with the United State Postal Service "Express Mail Post Office to Addressee" service under 37 CFR §1.10 on the date indicated above and is addressed to: BOX APPLICATION, Assistant Commissioner for Patents, Washington, DC 20231

Peter J. Manso December 22, 1999
Name of person mailing paper of fee Date


Signature

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re patent application of: Dave Bova
For: ***NICOTINIC ACID COMPOSITIONS FOR TREATING
HYPERLIPIDEMIA AND RELATED METHODS THEREFOR***
Examiner:
Art Unit:
Express Mail Label No.: EE683174198US

BOX APPLICATION
Assistant Commissioner
for Patents
Washington, D.C. 20231

Sir:

PRELIMINARY AMENDMENT

It is hereby respectfully requested that the above-identified application for U.S. patent be amended as follows.

In the Title:

In the title, please add - - HYDROPHOBIC COMPONENT FREE SUSTAINED RELEASE
- - before "NICOTINIC".

In the Specification:

On page 1, line 5, please delete "which is a continuation" and substitute the following therefore - - filed under U.S.C., Title 35, §120 and C.F.R., Title 37, §§1.53(b) and 1.78 and is continuation of U.S. Patent Application, Serial No. 08/814,974 filed on March 6, 1997, which is a continuation-in-part--


Remarks

It is respectfully submitted that all currently pending claims are patentably distinct over the disclosures of record and that the above-identified application for U.S. patent is in condition for allowance and early passage of this patent application to issuance is earnestly solicited.

Respectfully submitted.

AKERMAN SENTERFITT & EIDSON, P.A.

By


Peter J. Manso

Registration No. 32264

Akerman, Senterfitt & Eidson, P.A.
Las Olas Centre II, Suite 1600
350 East Las Olas Boulevard
Fort Lauderdale, Florida 33301
Telephone: 954-463-2700
Facsimile: 954-463-2224

NICOTINIC ACID COMPOSITIONS FOR TREATING HYPERLIPIDEMIA AND RELATED METHODS THEREFOR

Related Patent Applications

5 This application for U.S. patent is a U.S.C., Title 35, §111(a) application
which is a continuation of U.S. Patent Application, Serial No. 08/368,378 filed January
14, 1995, which is a continuation-in-part of U.S. Patent Application, Serial No.
08/124,392, filed September 20, 1993.

Field of the Invention

10 This invention generally relates to compositions of nicotinic acid useful for
treating hyperlipidemia and methods of treating hyperlipidemia employing such
compositions. More particularly, the present invention employs a composition of
nicotinic acid, derivatives and mixtures thereof, and a swelling agent to form a time
release sustaining composition for nocturnal or evening dosing. Specifically, the
present invention employs a composition of nicotinic acid and hydroxypropyl
15 methylcellulose to treat hyperlipidemia in a once per day oral dosage form given during
the evening hours.

Background

Nicotinic acid has been used for many years in the treatment of hyperlipidemia.
This compound has long been known to exhibit the beneficial effects of reducing total

cholesterol, low density lipoproteins or "LDL cholesterol", triglycerides and apolipoprotein a (Lp(a)) in the human body, while increasing desirable high density lipoproteins or "HDL cholesterol".

Nicotinic acid has normally been administered three times per day after meals.

5 This dosing regimen is known to provide a very beneficial effect on blood lipids as discussed in Knopp et al; "Contrasting Effects of Unmodified and Time-Release Forms of Niacin on Lipoproteins in Hyperlipidemic Subjects: Clues to Mechanism of Action of Niacin"; Metabolism 34/7, 1985, page 647. The chief advantage of this profile is the ability of nicotinic acid to decrease total cholesterol, LDL cholesterol, triglycerides
10 and Lp(a) while increasing HDL particles. While such a regimen does produce beneficial effects, cutaneous flushing and the like still often occurs in the hyperlipidemics to whom the compound is administered.

In order to avoid or reduce the cutaneous flushing, a number of materials have been suggested for administration with an effective antihyperlipidemic amount of
15 nicotinic acid, including guar gum in U.S. Pat. No. 4,965,252, and mineral salts as disclosed in U.S. Pat. No. 5,023,245; or inorganic magnesium salts as reported in U.S. Pat. No. 4,911,917. These materials have been reported to avoid or reduce the cutaneous flushing side effect commonly associated with nicotinic acid treatment.

Another method of avoiding or reducing the side effects associated with
20 immediate release niacin is the use of sustained release formulations. Sustained release formulations are designed to slowly release the compound from the tablet or capsule. The slow drug release reduces and prolongs blood levels of drug and thus minimizes the side effects. Sustained release formulations of niacin have been developed, such as Nicobid™ capsules (Rhone-Poulenc Rorer), Endur-acin™ (Innovite Corporation) and Pat.

No. 5,126,145 which describes a sustained release niacin formulation containing two different types of hydroxypropyl methylcellulose and a hydrophobic component.

Studies in hyperlipidemic patients have been conducted with a number of sustained release niacin products. These studies have demonstrated that the sustained release products do not have the same advantageous lipid altering effects as immediate release niacin, and in fact often have a worse side effect profile compared to the immediate release product. The major disadvantage of the sustained release formulations, as can be seen in Knopp et al., 1985, is the significantly lower reduction in triglycerides (-2% for the sustained release versus - 38% for the immediate release) and lower increase in HDL cholesterol, represented as HDL₂ particles which are known by the art to be most beneficial, (-5% for the sustained release versus +37% for the immediate release).

Additionally, sustained release niacin formulations have been noted as causing greater incidences of liver toxicity as described in Henken et al (Am J Med 91:1991 1991) and Dalton et al (Am J Med 93: 102 1992). There is also great concern regarding the potential of these formulations in disrupting glucose metabolism and uric acid levels.

In a recent edition of the JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION (JAMA), an article appeared which presented research results investigating the liver toxicity problems associated with a sustained release form of nicotinic acid. "A Comparison of the Efficacy and Toxic Effects of Sustained- vs. Immediate-Release Niacin in Hypercholesterolemic Patients", McKenney et al., JAMA, Vol. 271, No. 9, March 2, 1994, page 672. The article presented a study of twenty-three patients. Of that number, 18 or 78 percent were forced to withdraw because liver function tests (LFTs) increased indicating potential liver damage. The conclusion of

the authors of that article was that the sustained release form of niacin "should be restricted from use."

5 A similar conclusion was reached in an article authored by representatives of the Food and Drug Administration and entitled "Hepatic Toxicity of Unmodified and Time-Release Preparations of Niacin", Rader, et al., THE AMERICAN JOURNAL OF MEDICINE, Vol. 92, January 1992, page 77. Because of these studies and similar conclusions drawn by other health care professionals, the sustained release forms of niacin have experienced limited utilization.

10 Therefore, it can be seen from the scientific literature that there is a need for development of a sustained release niacin formulation and a method of delivering said formulation which would provide hyperlipidemic patients with "balanced lipid alteration", i.e. reductions in total cholesterol, LDL cholesterol, triglycerides and Lp(a) as well as increases in HDL particles, with an acceptable safety profile, especially as regards liver toxicity and effects on glucose metabolism and uric acid levels.

15 **Summary of the Invention**

In brief, the present invention alleviates and overcomes certain of the above-identified problems and shortcomings of the present state of nicotinic acid therapy through the discovery of novel nicotinic acid formulations and methods of treatment.

20 It is therefore, an object of the present invention to provide a composition of nicotinic acid or any compound which is metabolized by the body to form nicotinic acid for treating hyperlipidemia.

It is another object of the present invention to provide a composition as above, which has a time release sustaining characteristic.

It is yet another object of the present invention to provide a method for employing a composition as above, for treating hyperlipidemia, which results in little or no liver damage.

At least one or more of the foregoing objects, together with the advantages thereof over the known art relating to the treatment of hyperlipidemia, which shall become apparent from the specification which follows, are accomplished by the invention as hereinafter described and claimed.

In general the present invention provides an improved antihyperlipidemia composition of the oral type employing an effective antihyperlipidemic amount of nicotinic acid, wherein the improvement comprises compounding the nicotinic acid with from about 5% to about 50% parts by weight of hydroxypropyl methylcellulose per hundred parts by weight of tablet or formulation.

The present invention also provides an orally administered antihyperlipidemia composition which comprises from about 30% to about 90% parts by weight of nicotinic acid; and, from about 5% to about 50% parts by weight of hydroxypropyl methylcellulose.

The present invention also includes a method of treating hyperlipidemia in a hyperlipidemic. The method comprises the steps of forming a composition which comprises an effective antihyperlipidemic amount of nicotinic acid and an amount of excipients to provide sustained release of drug. The method also includes the step of orally administering the composition to the hyperlipidemic nocturnally.

A method of treating hyperlipidemia in a hyperlipidemic according to the invention, comprises dosing the hyperlipidemic with an effective antihyperlipidemic amount of nicotinic acid or compound metabolized to nicotinic acid by the body. The dose is given once per day in the evening or at night, combined with a pharmaceutically

acceptable carrier to produce a significant reduction in total and LDL cholesterol as well as a significant reduction in triglycerides and Lp(a), with a significant increase in HDL cholesterol.

5 The above features and advantages of the present invention will be better understood with reference to the following detailed description and examples. It should also be understood that the particular methods and formulations illustrating the present invention are exemplary only and not to be regarded as limitations of the present invention.

Detailed Description of the Invention

10 By way of illustrating and providing a more complete appreciation of the present invention and many of the attendant advantages thereof, the following detailed description and examples are given concerning the novel methods and formulations.

15 The present invention employs nicotinic acid or a compound other than nicotinic acid itself which the body metabolizes into nicotinic acid, thus producing the same effect as described herein. The other compounds specifically include, but are not limited to the following: nicotinyl alcohol tartrate, d-glucitol hexanicotinate, aluminum nicotinate, niceritrol and d,1-alpha-tocopheryl nicotinate. Each such compound will be collectively referred to hereinbelow by "nicotinic acid."

20 As stated hereinabove, nicotinic acid has been employed in the past for the treatment of hyperlipidemia, which condition is characterized by the presence of excess fats such as cholesterol and triglycerides, in the blood stream. According to the present invention, a sustained release composition of nicotinic acid is prepared as an example. By "sustained release" it is understood to mean a composition which when orally administered to a patient to be treated, the active ingredient will be released for
25 absorption into the blood stream over a period of time. For example, it is preferred

that in a dosage of about 1500 milligrams (hereinafter "mgs") of nicotinic acid, approximately 100 percent of the nicotinic acid will be released to the blood stream in about 4 to about 24 hours.

The specific sustained release composition according to the present invention employs an effective antihyperlipidemic amount of nicotinic acid. By "effective antihyperlipidemic amount" it is understood to mean an amount which when orally administered to a patient to be treated, will have a beneficial effect upon the physiology of the patient, to include at least some lowering of total cholesterol, LDL cholesterol, triglycerides and Lp(a) and at least some increase in HDL cholesterol in the patient's blood stream. An exemplary effective antihyperlipidemic amount of nicotinic acid would be from about 250 mgs to about 3000 mgs of nicotinic acid to be administered according to the invention as will be more fully described hereinbelow. This amount will vary dependent upon a number of variables, including the psychological needs of the patient to be treated.

Preferably, there is also included in the sustained release composition according to the present invention, a swelling agent which is compounded with the nicotinic acid, such that when the composition is orally administered to the patient, the swelling agent will swell over time in the patient's gastrointestinal tract, and release the active nicotinic acid, or a compound which produces nicotinic acid into the gastrointestinal system for absorption into the blood stream, over a period of time. As is known in the art, such swelling agents and amounts thereof, may be preselected in order to control the time release of the active ingredient. Such swelling agents include, but are not limited to, polymers such as sodium carboxymethylcellulose and ethylcellulose and waxes such as bees wax and natural materials such as gums and gelatins or mixtures of any of the above. Because the amount of the swelling agent will vary depending upon the nature

of the agent, the time release needs of the patient and the like, it is preferred to employ amounts of the agent which will accomplish the objects of the invention.

An exemplary and preferred swelling agent is hydroxypropyl methylcellulose, in an amount ranging from about 5% to about 50% parts by weight per 100 parts by weight of tablet or formulation. The preferred example will ensure a sustained time release over a period of approximately 4-24 hours as demonstrated by in vitro dissolution techniques known to the art.

A binder may also be employed in the present compositions. While any known binding material is useful in the present invention, it is preferred to employ a material such as one or more of a group of polymers having the repeating unit of 1-ethenyl-2-pyrrolidinone. These polymers generally have molecular weights of between about 10,000 and 700,000, and are also known as "povidone".

Amounts of the binder material will of course, vary depending upon the nature of the binder and the amount of other ingredients of the composition. An exemplary amount of povidone in the present compositions would be from about 1% to about 5% by weight of povidone per 100 parts by weight of the total formulation.

Processing aids such as lubricants, including stearic acid, may also be employed, as is known in the art. An exemplary amount of stearic acid in the present compositions would be from about 0.5% to about 2.0% by weight per 100 parts by weight of tablet or formulation.

Examples of various embodiments of the present invention will now be further illustrated with reference to the following examples.

General Experimental

In order to demonstrate the effectiveness of the compositions and method of the present invention over known antihyperlipidemia compositions and methods heretofore

known in the art, a number of substantially identical composition were prepared according to the disclosure hereinabove. The composition ingredients and amounts are listed in TABLE IA hereinbelow.

TABLE IA
Test Tablet Composition

5

<u>Ingredient</u>	<u>375 mg.</u>	<u>500 mg</u>	<u>750 mg</u>
Nicotinic Acid	375.0	500.0	750.0
Hydroxypropyl methylcellulose	188.7	203.0	204.7
10 Povidone	12.9	17.2	25.9
Stearic Acid	5.8	7.3	9.9
TOTAL	582.4 mg	727.5 mg	990.5 mg

15

The ingredients were compounded together to form a tablet. More specifically, Niaspan® once-daily tablets in accordance with the present invention utilize a hydrophilic matrix controlled drug delivery system. This is a dynamic system composed of polymer wetting, polymer hydration and polymer disintegration/dissolution. The mechanism by which drug release is controlled depends on, for example, initial polymer wetting, expansion of the gel layer, tablet erosion and niacin solubility. After initial wetting, the hydrophilic polymer starts to partially hydrate, forming a gel layer. As water permeates into the tablet increasing the thickness of the gel layer, drug diffuses out of the gel layer. As the outer layer of the tablet becomes fully hydrated it erodes. It is believed that this erosion results in additional drug release. The controlled release from this matrix delivery system can be modified depending on the type and molecular weight of hydrophilic polymer used.

20

25

A Niaspan® formulation consists of Niacin, Methocel® E10M Premium, Povidone K90 and Hystrene 5016 (stearic acid). Methocel® E10M Premium is utilized as a controlled-release agent in the Niaspan® formulation. Methocel is a partly O-

methyated and O-(2-hydroxypropylated) cellulose and is available in several grades which vary in terms of viscosity and degree of substitution. Methocel is manufactured by Dow Chemical.

5 Povidone K90 is employed as a granulating/binding agent in a Niaspan® formulation. Povidone is a synthetic polymer consisting of linear 1-vinyl-2-pyrrolidone groups, the degree of polymerization of which results in polymers of various molecular weights, or as indicated above. It is characterized by its viscosity in aqueous solution, relative to that of water, expressed as a K-value, ranging from 10-120. Povidone K90 has an approximate molecular weight of 1,000,000. Povidone is a hygroscopic, water
10 soluble material. Povidone K90 present in a Niaspan® formulation is manufactured by ISP (International Specialty Products). Hystrene 5016 is utilized as an external lubricant in the Niaspan® formulation. Hystrene 5016 is a mixture of stearic acid and palmitic acid. The content of stearic acid is not less than about 40.0% and the sum of the two acids is not less than about 90.0%. Hystrene 5016 is manufactured by Witco.
15 Refer to Table IB for Niaspan® formulation details.

Qualitatively, the four tablet strength formulations are identical. The major component of each formulation is a granulated mixture of Niacin, Methocel E10M and Povidone K90. The granulation process improves compression properties.

TABLE IB
Niaspan® Tablet Formulations

Niaspan® Product	375mg Tablets	500mg Tablets	750mg Tablets	1000mg Tablets
<u>Formulation, %/Tablet</u>				
Niacin	64.4	70.5	77.4	83.1
Methocel E10M	7.4	8.1	8.9	9.5
Premium (Intrgranular)				
Povidone K90	2.2	2.4	2.7	2.9
Methocel E10M				
Premium (Extragranular)	25.0	18.0	10.0	3.5
Hystrene 5016 (Stearic Acid)	1.0	1.0	1.0	1.0
Tablet weight, mg	582.5	709.5	968.6	1203.6

Niaspan® formulations are presented in white caplet shape tablets. Caplet dimensions differ with respect to product strength. The 375mg and 500mg Niaspan® tablets are compressed with tooling measuring approximately 0.687" in length x 0.281" by width. The length and width of the 750mg and 1000mg tooling measures approximately 0.750" x 0.320". Target tablet weight and hardness dictate thickness across the four Niaspan® products. The production of the Niaspan® tablets will now be described generally as set forth below.

Niaspan® Granulation Process Flow Chart

	<u>Raw Materials</u>	<u>Process Flow</u>	<u>Equipment</u>
5	Niacin Povidone K90 Methocel E10M (Intragranular) Purified Water	Granulate	High shear granulator (Littleford FM130)
		↓	
		Dry	Fluid bed drier (Glatt fluid bed drier)
		↓	
		Parcel size reduction	Mill (Kemutec Betagrind)

Niaspan® Granulation Process Description

10 Niaspan® granulation raw materials are dispensed and granulated in a high shear granulator. The wet granules are sieved into a fluid bed drier and are dried. When the drying process is complete, the granules are milled. Milling ensures uniform particle size distribution throughout the Niaspan® granulation.

Niaspan® Tablet Process Flow Chart

	<u>Raw Materials</u>	<u>Process Flow</u>	<u>Equipment</u>
		<u>Niaspan® Tablet Blend</u>	
5	Methocel E10M (Extragranular Hystrene 5016 (Stearic acid)	Blend Milled Niaspan® granules with extragranular Methocel E10M and Hystrene 5016	Blender (Patterson-Kelley V- Blender)
		↓	
		<u>Niaspan® Tablet Manufacture</u>	
		Compress Niaspan® Tablet Blend	Rotary tablet press

Niaspan® Tablet Process Description

10 A Niaspan® tablet blend is manufactured by blending the Niaspan® granulation, extragranular Methocel E10M and Hystrene 5016. The quantities of each Niaspan® tablet blend component will depend on the particular Niaspan® dose being manufactured (refer to Table IB). A Niaspan® tablet blend is compressed to form Niaspan® tablets. Niaspan® tablet physical properties will vary depending on the particular Niaspan® dose
15 being manufactured.

 Production of Niaspan® tablets will now be discussed in greater detail. The initial stage of manufacturing is the same for all four tablet strengths of Niaspan® (375,500, 750, and 1000mg). One batch of Niaspan® granulation is comprised of four individual 40.0kg units of granulation which are processed separately, but under like
20 conditions. The four individual granulations are sampled and tested individually and subsequently released for blending. The base granulation is not strength specific and may be used to manufacture any tablet strength of Niaspan®.

 The ingredients in the base granulation are set forth in Table IC below:

TABLE IC

Component	Function	Quantity per kilogram granulation (kg)	% per kilogram granulation (%)	Quantity per 160.00 kg batch (kg)
Niacin, USP	Drug Substance	0.87	87.00	139.20
Povidone, USP	Binder	0.03	3.00	4.80
Methocel USP, E10M Premium CR Grade	Controlled-Release Agent	0.10	10.00	16.00
Purified Water, USP*	Granulation Reagent	0.00*	0.00*	48.00
Total				160.00

*Purified Water, USP is used as a granulation reagent and does not appear in the finished granulation.

Raw materials are quantitatively dispensed into appropriately labeled double polyethylene-lined containers using calibrated scales. Purified Water, USP is dispensed into an appropriate vessel from which it is later pumped during the wet-massing operation.

A Littleford FM130 granulator is charged with approximately one half of the Niacin, USP required for the process unit (~ 17.4 kg) followed by about 4.00kg of Methocel, USP E10M Premium CR Grade; about 1.20kg of Povidone, USP; and the balance of the Niacin, SP (~ 17.40kg). The powder bed is dry mixed in the Littleford FM130 granulator, with choppers on, for approximately 1 minute. At the completion of the 1-minute pre-mix cycle, about 12.0 ± 0.05 kg of Purified Water, USP are sprayed onto the powder bed at a rate of about 2.40 ± 0.24 kg/minute. Immediately following the addition of the Purified Water, USP, the unit is granulated for about 5 minutes.

The granulated unit is discharged into double polyethylene-lined containers and then manually loaded into a Glatt bowl while being passed through a #4 mesh screen. the Glatt bowl is loaded into a Glatt TFO-60 fluid-bed drier with an inlet air

temperature setting of about $70^{\circ}\text{C} \pm 5^{\circ}\text{C}$. The unit is dried until a moisture level of $\leq 1.0\%$ is obtained as determined using a Computrac® Moisture Analyzer, model MA5A. The dried granulation is discharged into appropriately labeled, double polyethylene-lined drums and reconciled.

5 The dried and reconciled granulation is passed through a Kemutec BetaGrind mill equipped with a 1.5mm screen and running at approximately 1500 RPM. The milled granulation is collected into appropriately labeled, double polyethylene-lined drums and reconciled. The milled granulation is sampled and tested by Quality Control and released prior to further processing.

10 The released granulation units are charged to a Patterson-Kelley 20 ft³ V-blender after which they are blended together for about 10 ± 1 minutes and then discharged to appropriately labeled, double polyethylene-lined containers.

As stated above, Niaspan® tablets are formulated from a common granulation which is blended with appropriate quantities of Methocel, USP E10M Premium CR
15 Grade and Stearic Acid, NF to achieve the final dosage formulation. Tables IA and IB describe the formulation for each Niaspan® tablet strength, 375mg, 500mg, 750mg, and 1000mg, respectively.

Two study groups consisting of eleven and fourteen patients each were formed. Blood samples were taken from the patients, and tested for total cholesterol, LDL
20 cholesterol, triglycerides and HDL cholesterol to establish baseline levels from which fluctuations in these lipids could be compared. The patients were then placed upon a regimen of the above discussed tablets, totalling approximately 1500 mg of nicotinic acid, once per day before going to bed. After eight weeks of this regimen, the patients were again tested for lipid profiles. The results of the tests conducted at eight weeks,
25 showing the changes in the lipid profiles as a percentage change from the baseline, are

reported in the table hereinbelow. Positive numbers reflect percentage increases and negative numbers reflect percentage decreases in this table.

TABLE II
Patient Study Lipid Profile Data

	<u>Pt. No.</u>	<u>Total-C</u>	<u>LDL-C</u>	<u>Apo B</u>	<u>Trigs</u>	<u>HDL-C</u>	<u>HDL₂-C</u>	<u>Lp(a)</u>
5	GROUP A							
	1	-8.2	-12.0	NA	-17.3	22.0	NA	NA
	2	-5.9	-27.0	NA	-28.7	65.0	NA	NA
	3	-15.1	-13.0	NA	-22.0	-9.1	NA	NA
10	4	-3.3	-10.0	NA	61.6	3.8	NA	NA
	5	-16.5	-17.7	NA	-28.8	11.1	NA	NA
	6	-12.4	-25.9	NA	-42.0	51.6	NA	NA
	7	-24.2	-31.4	NA	-39.4	12.5	NA	NA
	8	-6.7	-7.4	NA	-42.4	18.8	NA	NA
15	9	4.5	1.1	NA	7.2	9.2	NA	NA
	10	2.8	-0.2	NA	-2.7	22.9	NA	NA
	11	-13.0	-9.4	NA	-54.0	44.3	NA	NA
	Mean	-8.9	-13.9	NA	-18.9	23.0	NA	NA
	p-Value	0.0004	0.0001		0.0371	0.0068		
20	GROUP B							
	1	-19.2	-27.1	-24.4	-33.4	20.0	22.3	-81.9
	2	-32.2	-35.7	-28.0	-60.4	4.3	3.2	-25.3
	3	-21.4	-33.6	-35.6	-33.4	30.4	38.6	-17.4
	4	-19.9	-24.6	-15.1	-20.8	9.6	16.1	-27.0
25	5	-3.3	-2.1	-29.4	-41.1	5.8	2.4	-22.4
	6	PATIENT WITHDREW FROM STUDY						
	7	23.1	-32.6	-42.6	-58.6	49.2	68.9	-14.3
	8	24.8	34.0	-28.4	5.5	6.5	-6.8	NA

TABLE II (Continued)

<u>Pt. No.</u>	<u>Total-C</u>	<u>LDL-C</u>	<u>Apo B</u>	<u>Trigs</u>	<u>HDL-C</u>	<u>HDL-C</u> <u>C</u>	<u>Lp(a)</u>
GROUP B							
9	10.1	12.0	-16.8	-11.6	20.7	-12.3	40.6
10	-2.9	-7.7	-28.0	-59.0	53.1	70.5	-41.2
11	-10.5	-18.8	-25.3	-53.4	31.8	39.7	NA
5 12	-20.0	-30.8	-30.4	11.7	21.1	25.0	-28.4
13	17.4	16.8	-17.5	-17.5	51.3	51.9	38.5
14	-9.4	-16.6	-32.0	-46.9	52.3	67.6	17.6
Mean	-8.7	-12.8	-32.2	-27.2	25.3	30.1	-17.9
p-Value	0.0002	<0.0001	0.0001	<0.001	<0.0001	0.0002	<0.0188
10 Combined	-8.7	-13.3	Gp B	-26.1	25.3	Gp B	Gp B
p-Value	0.0002	<0.0001	only	<.0001	<0.0001	only	only

The data reported in TABLE II shows that the LDL levels in the Group A patients had a mean decrease of -13.9% and triglyceride decrease of -18.9% HDL cholesterol levels, the beneficial cholesterol, were raised by 23.0% in this Group. Similar results were obtained with the Group B patients. These studies demonstrate that dosing the sustained release formulation during the evening hours or at night provides reductions in LDL cholesterol levels equal to immediate release niacin on a milligram per milligram basis, but superior reductions in triglyceride reductions when compared to sustained release formulations dosed during daytime hours on a milligram per milligram basis. Additionally, the increases in HDL cholesterol obtained from dosing the sustained release formulation during the evening or at night were +23.0% for one group and +25.3% for the other group. Dosing during the evening therefore provides reduction in LDL cholesterol plus significant decreases in triglycerides and increases in HDL cholesterol with once-a-day dosing.

Groups A and B were also tested for liver enzymes (AST, ALT and Alkaline Phosphatase), uric acid and fasting glucose levels at the start of the study described

hereinabove (to form a baseline) and at two, four and eight week intervals. The results of these tests are listed in TABLES III-VII hereinbelow.

TABLE III
THE EFFECT OF NIASPAN® THERAPY ON AST (SGOT) LEVELS (U/L)
(1500 mgs dosed once-a-day at night)
(n = 28)

Weeks of Therapy With NIASPAN™

	<u>Pt #</u>	<u>Baseline</u>	<u>2 Wks.</u>	<u>4 Wks.</u>	<u>8 Wks.</u>	<u>Reference Range</u>
	GROUP A					
10	1	28	29	25	24	0-50
	2	24	25	24	26	0-50
	3	17	18	22	21	0-50
	4	14	16	15	17	0-50
	5	22	NA	32	52	0-50
15	6	21	17	17	14	0-50
	7	17	17	14	18	0-50
	8	20	21	22	22	0-50
	9	16	16	17	20	0-50
	10	18	21	21	25	0-50
20	11	21	21	22	21	0-50
	GROUP B					
	1	23	25	38	33	0-50
	2	20	20	21	21	0-50
	3	15	20	18	19	0-50
25	4	25	22	25	26	0-50
	5	23	21	17	18	0-50
	6	PATIENT WITHDREW DUE TO FLUSHING				
	7	21	18	18	19	0-50
	8	18	19	18	19	0-50
30	9	15	16	18	15	0-50
	10	16	15	19	28	0-50

TABLE III (Continued)

	<u>Pt #</u>	<u>Baseline</u>	<u>2 Wks.</u>	<u>4 Wks.</u>	<u>8 Wks.</u>	<u>Reference Range</u>
	GROUP B					
	11	20	22	24	28	0-50
	12	23	25	28	22	0-50
	13	20	15	20	19	0-50
5	14	18	25	20	18	0-50
	Combined Mean	19.8	20.4	20.8	21.1	
	Change From Baseline		+3.0%	+5.1%	+6.6%	
10	Level of Significance: p=0.4141					

TABLE IV
THE EFFECT OF NIASPAN® THERAPY ON ALT (SGPT) LEVELS (U/L)
(1500 mgs dosed once-a-day at night)
(n = 28)

Weeks Of Therapy With NIASPAN®

	<u>Pt#</u>	<u>Baseline</u>	<u>2 Wks.</u>	<u>4 Wks.</u>	<u>8 Wks.</u>	<u>Reference Range</u>
	GROUP A					
	1	32	28	39	30	0-55
20	2	24	25	23	26	0-55
	3	18	23	30	30	0-55
	4	7	13	14	14	0-55
	5	14	NA	43	46	0-55
	6	22	11	14	10	0-55
25	7	9	7	11	7	0-55
	8	16	18	23	21	0-55
	9	14	17	20	14	0-55
	10	14	15	17	19	0-55
	11	18	18	20	16	0-55

00470603 42299
00001 00902450

TABLE IV (Continued)

	<u>Pt#</u>	<u>Baseline</u>	<u>2 Wks.</u>	<u>4 Wks.</u>	<u>8 Wks.</u>	<u>Reference Range</u>
	GROUP B					
	1	16	17	27	29	0-55
	2	16	14	15	22	0-55
	3	13	21	13	16	0-55
5	4	23	20	26	17	0-55
	5	21	23	17	15	0-55
	6	PATIENT WITHDREW DUE TO FLUSHING				
	7	21	16	18	21	0-55
	8	18	20	17	18	0-55
10	9	11	5	11	8	0-55
	10	8	10	14	17	0-55
	11	17	12	18	16	0-55
	12	14	18	20	16	0-55
	13	14	NA	11	10	0-55
15	14	23	23	19	19	0-55
	Combined Mean	17.7	17.5	19.3	18.2	
	Change From Baseline		-1.1%	9.0%	+2.8%	
20	Level of Significance: p=0.3424					

Downloaded from www.ascp.com

5

20

	<u>Pt#</u>	<u>Baseline</u>	<u>2 Wks.</u>	<u>4 Wks.</u>	<u>8 Wks.</u>	<u>Reference Range</u>
	GROUP A					
10	1	52	56	57	55	20-140
	2	103	100	89	102	20-140
	3	54	45	53	51	20-140
	4	70	68	71	91	20-140
	5	77	NA	74	81	20-140
15	6	55	48	49	51	20-140
	7	72	71	79	75	20-140
	8	55	49	47	50	20-140
	9	53	55	56	45	20-140
	10	74	73	75	75	20-140
	11	18	18	20	16	20-140
20	GROUP B					
25	1	73	67	89	95	20-140
	2	82	64	72	71	20-140
	3	73	69	72	82	20-140
	4	37	36	37	38	20-140
	5	65	53	54	61	20-140
	6	PATIENT WITHDREW DUE TO FLUSHING				
	7	64	58	58	58	20-140
	8	79	78	65	73	20-140
	9	94	92	103	93	20-140
	10	69	67	70	65	20-140
	11	59	67	63	72	20-140
	12	65	59	59	63	20-140
30						

TABLE V (Continued)

	<u>Pt#</u>	<u>Baseline</u>	<u>2 Wks.</u>	<u>4 Wks.</u>	<u>8 Wks.</u>	<u>Reference Range</u>
	GROUP B					
	13	64	68	66	64	20-140
	14	72	61	59	64	20-140
5	Combined Mean	66.5	61.5	63.3	65.8	
	Change From Baseline		-6.1%	-3.4%	+0.005%	
	Level of Significance: $p=0.0236$					

TABLE VI
THE EFFECT OF NIASPAN® THERAPY ON URIC ACID LEVELS (mg/dL)
(1500 mgs dosed once-a-day at night)
(n = 28)

5

Weeks Of Therapy With NIASPAN®

	<u>Pt#</u>	<u>Baseline</u>	<u>2 Wks.</u>	<u>4 Wks.</u>	<u>8 Wks.</u>	<u>Reference Range</u>
	GROUP A					
10	1	5.2	5.0	4.8	4.3	4.0-8.5
	2	4.0	4.6	4.5	6.2	2.5-7.5
	3	6.3	7.0	6.5	6.2	4.0-8.5
	4	3.1	4.6	4.2	3.8	2.5-7.5
	5	3.4	NA	3.3	4.2	2.5-7.5
	6	6.6	5.5	5.6	4.7	4.0-8.5
15	7	3.8	4.5	4.3	4.9	2.5-7.5
	8	4.4	3.8	5.1	4.5	2.5-7.5
	9	3.9	4.5	4.6	3.5	2.5-7.5
	10	2.6	2.9	2.8	2.7	2.5-7.5
	11	4.7	5.5	5.2	5.3	2.5-7.5
	GROUP B					
20	1	3.7	4.2	4.7	3.5	2.5-7.5
	2	2.8	3.5	3.6	2.3	4.0-8.5
	3	4.2	5.3	5.5	5.3	2.5-7.5
	4	4.7	3.9	5.1	3.6	4.0-8.5
	5	3.7	4.1	4.1	3.8	2.5-7.5
25	6	PATIENT WITHDREW DUE TO FLUSHING				
30	7	5.8	6.6	6.6	6.8	2.5-7.5
	8	4.7	4.3	5.4	5.6	2.5-7.5
	9	3.7	4.6	5.1	3.8	2.5-7.5
	10	4.2	5.0	4.4	8.5	2.5-7.5
	11	1.9	3.0	2.8	5.0	2.5-7.5
	12	5.6	5.4	6.2	5.6	4.0-8.5
	13	4.2	4.6	4.6	5.3	2.5-7.5

TABLE VI (Continued)

	<u>Pt#</u>	<u>Baseline</u>	<u>2 Wks.</u>	<u>4 Wks.</u>	<u>8 Wks.</u>	<u>Reference Range</u>
	GROUP B					
	14	5.5	5.4	6.1	5.3	2.5-7.5
	Combined Mean	4.54	4.82	4.92	4.86	*p=0.3450
5	Change From Baseline		+6.2%	+8.4%	+7.0%	

*Level of Significance: p=0.3450

Downloaded from www.ascp.com

TABLE VII
THE EFFECT OF NIASPAN® THERAPY
ON FASTING GLUCOSE LEVELS (mg/dL)
(1500 mgs dosed once-a-day at night)
(n = 28)

5

Weeks Of Therapy With NIASPAN®

	<u>Pt #</u>	<u>Baseline</u>	<u>2 Wks.</u>	<u>4 Wks.</u>	<u>8 Wks.</u>	<u>Reference Range</u>
	GROUP A					
10	1	114	122	123	110	70-115
	2	101	105	107	101	80-125
	3	99	98	109	103	70-115
	4	100	118	94	94	80-125
	5	89	NA	82	103	80-125
15	6	97	103	94	107	70-115
	7	85	107	100	94	80-125
	8	98	107	103	101	80-125
	9	97	97	100	110	80-125
	10	94	101	111	97	70-115
	11	102	103	95	95	80-125
20	GROUP B					
25	1	101	97	83	99	70-115
	2	90	95	96	89	80-125
	3	96	98	95	97	70-115
	4	116	139	113	125	80-125
	5	88	92	91	95	70-115
30	6	PATIENT WITHDREW DUE TO FLUSHING				
	7	106	114	118	117	70-115
	8	95	106	106	108	70-115
	9	81	92	84	92	70-115
	10	108	117	122	105	70-115
	11	85	106	106	108	70-115
	12	92	89	101	86	80-125
	13	99	105	94	100	70-125
	14	100	108	84	107	70-125

TABLE VII Continued

<u>Pt #</u>	<u>Baseline</u>	<u>2 Wks.</u>	<u>4 Wks.</u>	<u>8 Wks.</u>	<u>Reference Range</u>
Combined Mean	98.4	105.8	101.6	102.3	
Change From Baseline		+7.5%	+3.3%	+4.0%	

5 Level of Significance: $p=0.0021$

In order to provide a comparison between the state of the art prior to the present invention, and in order to quantify the magnitude of the improvement that the invention provides over the prior art, another study was conducted. This study included 240 patients dosed according to the present invention as described hereinabove.

10 Compared to this group was the group of patients studied by McKenney et al., as reported hereinabove. The results of this study are reported in TABLE VIII hereinbelow.

TABLE VIII
A Comparison of Changes in Liver Function Tests

		0	500	1000	1500	2000	2500	3000	TOTAL
5	McKenney SR^b Niacin^a								
	AST	23.8	27.9	40.4	36.6	56.5	na	97.0	
	%	--	117	170	154	237	na	408	
	Invention Dosage^c								
	AST	24.3	na	23.7	27.5	26.6	27.6	27.8	
	%	--	na	98	113	109	114	114	
10	McKenney SR Niacin								
	ALT	25.6	29.5	36.3	39.0	59.1	na	100.0	
	%	--	115	142	152	231	na	391	
	Invention Dosage								
	ALT	21.4	na	18.7	22.6	21.3	22.4	21.8	
	%	--	na	87	106	100	105	102	
15	McKenney SR Niacin								
	ALK	95	95	106	105	136	na	135	
	%	--	100	112	111	143	na	142	

TABLE VIII (continued)

		0	500	1000	1500	2000	2500	3000	TOTAL
Invention Dosage									
5	ALK	74.7	na	73.9	76.1	73.4	76.7	78.0	
	%	--	na	99	102	98	103	104	
	McKenney SR Niacin								
	Drop	--	0	2	2	7	na	7	18
	n	--	--	--	--	--	--	--	23
10	%	--	0	9	9	30	na	30	78
	Invention Dosage								
	Drop	--	--	0	0	0	0	0	0
	n	--	--	26	67	97	35	15	240
	%	--	--	0	0	0	0	0	0
15	1 year	--	--	15	46	77	31	15	184
	1 year	--	--	58	69	79	89	100	77
	Dosed twice-per-day as described in "A Comparison of the Efficacy and Toxic Effects of Sustained - vs Immediate - Release Niacin in Hypercholesterolemic Patients" by McKenney et al. <u>Journal of the American Medical Association</u> , March 2, 1994; Vol. 271, No. 9, pages 672-677.								
	^b SR is "sustained release"								
	^c Dosed once-per-day at night								

^a Dosed twice-per-day as described in "A Comparison of the Efficacy and Toxic Effects of Sustained - vs Immediate - Release Niacin in Hypercholesterolemic Patients" by McKenney et al. Journal of the American Medical Association, March 2, 1994; Vol. 271, No. 9, pages 672-677.

^b SR is "sustained release"

^c Dosed once-per-day at night

The results of the comparison of the studies reported in TABLE VIII show that the control group (the McKenney group) had 18 of 23, or 78 percent of the patients therein drop out of the test because of an increase in their respective liver function tests. The patients withdrew at the direction of the investigator. In comparison, a group of 240 patients treated according to the present invention had zero patients drop out, based upon the same criteria for withdrawal. The tests results reported above indicate that this sustained release dosage form caused no elevation in liver function tests (i.e., no liver damage), no elevations in uric acid and only a small, 7.5% increase in fasting glucose levels which in fact decreased during continued therapy.

Thus it should be evident that the compositions and method of the present invention are highly effective in controlling hyperlipidemia in hyperlipidemics, by reducing the levels of LDL cholesterol, triglyceride and Lp(a) while increasing HDL cholesterol levels. The present invention is also demonstrated not to cause elevations in liver function tests, uric acid or glucose levels for the hyperlipidemics.

Based upon the foregoing disclosure, it should now be apparent that the use of the compositions and methods described herein will carry out the objects set forth hereinabove. It is, therefore, to be understood that any variations in sustained release formulation evident fall within the scope of the claimed invention and thus, the selection of specific component elements can be determined without departing from the spirit of the invention herein disclosed and described. In particular, sustained release excipients, binders and processing aids according to the present invention are not necessarily limited to those exemplified hereinabove. Thus, the scope of the invention shall include all modifications and variations that may fall within the scope of the attached claims.

CLAIMS

What is claimed is:

1. A method of treating hyperlipidemia in a hyperlipidemic comprising dosing the hyperlipidemic with an effective antihyperlipidemic amount of nicotinic acid or compound metabolized to nicotinic acid by the body, once per day in the evening or at night combined with pharmaceutically acceptable carriers, to produce a reduction in total and LDL cholesterol, triglycerides and Lp(a), with a significant increase in HDL cholesterol.
2. A method, as set forth in Claim 1, wherein the hyperlipidemic is dosed with from about 250 parts to about 3000 parts by weight of nicotinic acid.
3. A method as set forth in Claim 1 which causes little or no serious liver damage, uric acid increases or elevations in fasting glucose levels.
4. A method as set forth in Claim 1 wherein the release rate of said nicotinic acid or compound metabolized by the body to nicotinic acid is from about 2.0% per hour to about 25% per hour.
5. A method as set forth in Claim 1 wherein said nicotinic acid or compound metabolized to nicotinic acid by the body is prepared by formulating the active compound with from about 5% to about 50% parts by weight of hydroxypropyl methylcellulose per 100 parts by weight of tablet.

6. A method, as set forth in Claim 1, wherein said nicotinic acid or compound metabolized to nicotinic acid by the body is dosed in the form of a sustained release formulation or tablet containing from about 1 to about 4 parts by weight of binder per 100 parts by weight of tablet.
- 5 7. A method, as set forth in Claim 4, wherein said binder is a polymer having the repeating polymerization unit 1-ethenyl-2-pyrrolidone.
8. A method, as set forth in Claim 1, wherein said nicotinic acid or compound metabolized to nicotinic acid by the body is dosed in the form of a sustained release formulation or tablet comprising from about 0.5 to about 2.5 parts by weight of a lubricating agent per 100 parts by weight of tablet.
- 10 9. A method, as set forth in Claim 8, wherein said lubricating agent is selected from the group consisting of stearic acid and magnesium stearate.
10. A method, as set forth in Claim 1, wherein the compound metabolized to nicotinic acid by the body to nicotinic acid is nicotinyl alcohol tartrate.
- 15 11. A method, as set forth in Claim 11, wherein the amount of nicotinyl alcohol tartrate is from about 100 milligrams to about 500 milligrams per dosage unit.
12. A method, as set forth in Claim 1, wherein the compound metabolized to nicotinic acid by the body is selected from the group consisting of: d-glucitol hexanicotinate, aluminum nicotinate, and, 1-alpha-tocopheryl nicotinate.

Abstract

An orally administered antihyperlipidemia composition according to the present invention includes from about 250 to about 3000 parts by weight of nicotinic acid, and from about 5 to about 50 parts by weight of hydroxypropyl methylcellulose. Also, a method of treating hyperlipidemia in a hyperlipidemic having a substantially periodic physiological loss of consciousness, includes the steps of forming a composition having an effective antihyperlipidemic amount of nicotinic acid and a time release sustaining amount of a swelling agent. The method also includes the step of orally administering the composition to the hyperlipidemic once per day "nocturnally," that is in the evening or at night.